



## Appendix to Biology Reports

### Body Fluid Identification

**Presumptive** - A non-confirmatory test used for detecting the possible presence of biological fluids. Presumptive tests make use of a target chemical to establish the possibility that a particular body fluid is present.

- Phenolphthalein is used as a presumptive test for blood.
- Hematrace is used as a presumptive test for human blood.
- Detection of acid phosphatase is used as a presumptive test for seminal protein.
- Detection of amylase is used as a presumptive test for saliva.
- Amylase may also be found in lower levels in urine, feces, perspiration, and vaginal secretions.

**Prostate Specific Antigen (PSA)** - A protein (also known as P30) produced by the prostate gland and found in semen. PSA concentration in semen is typically in levels far in excess of those found in other fluids.

**Spermatozoa** - The male reproductive cell that can be found in semen.

**Semen** is comprised of two components: the seminal plasma and spermatozoa. Seminal plasma contains PSA and acid phosphatase, typically in levels far in excess of those found in other fluids.

### Y-screening for Male DNA

The Y-screen assay is a rapid and reliable test to screen for male DNA in the processing of sexual assault kit samples. Utilizing a simplified extraction step which rapidly lyses cells (including sperm) and a quantification kit, the Y-screening rapidly detects the presence of male DNA in sexual assault kit swab evidence. The assay may be used in conjunction with, or independent from, other screening methods to assist in deciding whether to proceed with DNA analysis procedures.

### DNA Analysis

**DNA (Deoxyribonucleic Acid)**, the inherited genetic material found in most cells, contains markers that can differ from person to person. DNA analysis can determine these genetic markers and compare biological samples from different individuals.

**DNA Analysis** is comprised of several steps, including DNA extraction, DNA quantification, PCR/DNA amplification, and analysis of the resulting DNA alleles.

**Alleles** are an alternative form of DNA markers. Alleles are found at specific areas, or locations, of the DNA called **loci** (singular, **locus**).

**DNA extraction** recovers DNA from biological samples such as blood, saliva, semen, bone, hair, tissue, and skin cells.

**Differential Extraction** – A procedure in which sperm cells are separated from all other cells in a sample, resulting in a Sperm Fraction which is enriched for sperm DNA and a Non-Sperm/Epithelial Fraction which contains DNA from other cell types. Incomplete separation can occur and fractions may contain both sperm DNA and non-sperm DNA.



**Appendix to Biology Reports**

**DNA quantification** provides an estimate of the amount of DNA recovered from samples by using a technique called real time polymerase chain reaction (q-PCR). The quantification testing uses the **Applied Biosystems Quantifiler® Trio DNA Quantification Kit**. If sufficient DNA is detected, DNA amplification and analysis can be attempted.

The **PCR** (polymerase chain reaction) is a technique that copies specific areas of DNA. PCR generates large amounts of DNA from small starting amounts of DNA by repeated cycles of copying the DNA loci (**DNA amplification**); after amplification, the alleles present in the sample are identified.

PCR DNA testing for STRs uses a **DNA amplification kit**, a commercial product used to generate a DNA profile.

The MCCL uses the **Applied Biosystems AmpFISTR Globalfiler® PCR Amplification Kit** using 29 amplification cycles. Each STR locus tested in the Globalfiler® kit contains between 9 and 34 identifiable alleles. The loci tested are D3S1358, vWA, D16S539, CSF1PO, TPOX, D8S1179, D21S11, D18S51, D2S441, D19S433, TH01, FGA, D22S1045, D5S818, D13S317, D7S820, SE33, D10S1248, D1S1656, D12S391, D2S1338. The kit also tests the Amelogenin locus, Y-indel, and DYS391 which are used to determine the sex origin of a sample.

The MCCL uses the **Applied Biosystems AmpFISTR YFiler™ Plus PCR Amplification Kit** using 30 amplification cycles. Each STR locus tested in the YFiler® Plus kit contains between 8 and 23 identifiable alleles. The loci tested are DYS576, DYS389I, DYS635, DYS389II, DYS627, DYS460, DYS458, DYS19, GATAH4, DYS448, DYS391, DYS456, DYS390, DYS438, DYS392, DYS518, DYS570, DYS437, DYS385, DYS449, DYS393, DYS439, DYS481, DYF387S1, and DYS533.

**Stochastic effects** are defined as unequal sampling of the two alleles present from a heterozygous individual that result when only a few DNA copies are used to initiate PCR. Such samples may exhibit significantly different heterozygous peak heights or allelic dropout.

The MCCL uses a **stochastic threshold** of 400 RFUs for Globalfiler data as a means to ascertain if data detected is complete in mixtures and low-level samples. When a sample containing low-level amount of DNA template (0.10ng for single source samples) or a minor component(s) of a mixture exhibits peaks below **stochastic threshold**, it is possible that complete amplification has not occurred and not all alleles have been detected.

**STR** (short tandem repeat) loci contain alleles with a variable number of short repeating segments. Each STR allele can be described using a number that represents its number of repeats. A **DNA profile** is the collection of these numbers describing the DNA alleles found at an individual's STR DNA loci.

**Y-STR** loci contain alleles with a variable number of short repeating segments on the **Y chromosome**. Y-STRs are polymorphic among *unrelated* males and are inherited through the paternal line with little change through generations. Barring a mutation event, a person of interest's Y-STR haplotype will be the same in all paternal male relatives. Y-STRs differ from autosomal STRs in other ways. First, only male samples have a Y-STR haplotype because females do not possess a Y chromosome. Secondly, because the Y chromosome is inherited from the father, there is only a single allele at each locus, with the

Approved By	Gail M. Conklin	Revision	2021.1	Issued on	December 7, 2021
-------------	-----------------	----------	--------	-----------	------------------



**Appendix to Biology Reports**

exception of a duplication at DYS385 and/or DYF387S1. Lastly, the 27 loci amplified with the AmpFISTR Yfiler Plus® PCR Amplification kit are not independent of each other and represent a single haplotype.

Statistics

The rarity of a DNA profile can be expressed as an STR population frequency estimate, how often one would expect to see the DNA profile derived from the evidentiary item(s). The frequency estimate is expressed as a probability.

Autosomal STR population frequency estimates are based on the NIST population database (2013) and the National Research Council (1996) The Evaluation of Forensic DNA Evidence, Natl. Acad. Press, Washington DC.

The statistical values reported reflect the approximate frequency of the occurrence of a given DNA profile in a population of unrelated individuals. Therefore, these values may not be appropriate for relatives.

The Monroe County Crime Laboratory reports the most common frequency estimate of the three more common populations in the US: African-American, Caucasian, and Hispanic.

**Y-STR haplotype** - All Y-STR loci analyzed are physically linked on the Y-chromosome. Due to the lack of recombination, the entire Y-chromosome haplotype must be treated as a single locus. Haplotype frequencies are estimated using the **counting method**. The counting method involves searching a given haplotype against a database to determine the number of times the haplotype was observed in that database. The frequency of the haplotype in the database is then estimated by dividing the count by the number of haplotypes searched. The profile probability is estimated by applying a 95% confidence upper bound to the haplotype frequency, using the method described by Clopper and Pearson (1934) as per the SWGDAM Interpretation Guidelines for Y-Chromosome STR Typing by Forensic DNA Laboratories, 2014: Section 10.2.3.

The statistic reported is a count of the number of times that the haplotype in question was observed when searched against the database. The statistical strength of Y-STR testing is affected by the size of the population database searched. Profiles in the population database may not have results for all of the Y-STR loci. Therefore, the total number of observations in the database will vary dependent upon which loci are searched. Population databases may contain profiles typed with different multiplexes, containing different numbers and/or sets of loci, such that only a subset of the database may have been typed at all of the loci present in the evidence profile.

The MCCL uses the Y-Chromosome Haplotype Reference Database (<https://yhrd.org/>) for calculating profile probability. The Release Number of the Y-Chromosome Haplotype Reference Database is updated periodically. The statistical values may change based upon database size and composition. If prosecution of a case is anticipated, an updated calculation may be performed upon request.

Approved By	Gail M. Conklin	Revision	2021.1	Issued on	December 7, 2021
-------------	-----------------	----------	--------	-----------	------------------



**Appendix to Biology Reports**

Conclusions for DNA Comparisons

**The DNA profile obtained from Item X matches the DNA profile of Item Y:** When making a direct comparison between two specimens, a match may be reported if the DNA profile of the reference specimen is the same as the DNA profile of the questioned specimen(s).

**NOTE:** *If an evidentiary profile has been stated to match the known DNA profile of a specific individual, then the known DNA profiles of other individuals subsequently profiled in the case have been eliminated as possible sources of that specific evidentiary profile.*

**Item Y and Item Z cannot be excluded as a contributor(s) to the mixture:** For locations where comparisons could be made, all or most of the DNA alleles seen in an individual's DNA profile were represented in the mixture. The allele(s) that are not represented could be explained by any of several factors. Therefore, an individual cannot be ruled out as a possible contributor to the mixture.

**Item Y and Item Z are excluded as a contributor(s) to the mixture:** For locations where comparisons could be made, the DNA alleles seen in an individual's DNA profile were not represented in the mixture. The allele(s) that are not represented could not be explained. Therefore, an individual can be ruled out as a possible contributor to the mixture.

**The mixture is uninterpretable or inconclusive:** The DNA results from the evidence are either limited or too complex and are not suitable for comparison. Therefore, it cannot be determined whether an individual is a contributor to this mixture.

**Not suitable for comparison:** When a sample is amplified, DNA results may be inconclusive due to insufficient data or genetic complexity. Comparisons to known items and/or statistical calculations cannot be made. No further conclusions can be drawn regarding the source or sources of the resulting data.

CODIS

The main function of the **Combined DNA Index System (CODIS)** is to provide associations between cases and offenders or between two cases. All eligible DNA profiles that are suitable are entered into CODIS into the Local DNA Index System (LDIS). DNA profiles are searched a minimum of twice per week at LDIS. If a sample meets the completeness requirements for the state level, the profile is uploaded to the State DNA Index System (SDIS). If the DNA profile also meets the completeness requirements for the national level, the profile is uploaded to the National DNA Index System (NDIS). SDIS and NDIS searches are performed daily, Monday through Friday. NDIS may also perform additional searches. In the event of a new, positive association, a letter is generated by the MCCL notifying the investigating agency and their legal representative of any pertinent CODIS match information.

The notification letter will indicate the stringency of the association. A high stringency match indicates that all the alleles in the forensic specimen are the same as the candidate specimen. A moderate stringency match indicates that at least one locus has a different number of alleles between the forensic specimen and the candidate specimen, but all of the allelic values in the specimen with the fewest number of alleles are represented in the other specimen. A moderate stringency match may occur when one or both of the specimen profiles is a partial profile or a DNA mixture.

Approved By	Gail M. Conklin	Revision	2021.1	Issued on	December 7, 2021
-------------	-----------------	----------	--------	-----------	------------------



**Appendix to Biology Reports**

Conclusions for CODIS

**The DNA profile obtained from Item X was entered into the Local DNA Index System (LDIS) for comparison to forensic profiles. LDIS does not contain offender profiles.** The DNA profile does not meet the completeness requirements to be uploaded to the state or national level. The DNA profile will be searched at the local level only.

**The DNA profile obtained from Item X will also be uploaded to the State DNA Index System (SDIS) for comparison to forensic and offender profiles.** The DNA profile does not meet the completeness requirements to be uploaded to the national level. The DNA profile will be searched at the local and state levels.

**The DNA profile obtained from Item X will be uploaded to the State DNA Index System (SDIS) and National DNA Index System (NDIS) for comparison to forensic and offender profiles.** The DNA profile will be searched at the local, state, and national levels of CODIS.

***The DNA profile(s) obtained from Item(s) is/are not currently eligible for entry into the CODIS database for comparison purposes:*** Based on the current documentation in the laboratory case file, the documentation is lacking for one or more of the following general requirements:

- 1) That the item is related to a crime.
- 2) How the item is connected to the crime scene.
- 3) How the item (or DNA profile) is connected to the perpetrator of the crime.
- 4) That the item is **not** from a place where the perpetrator's profile can reasonably be expected to be found.

The laboratory can re-evaluate CODIS eligibility if more case documentation is submitted.

***The DNA profile obtained from Item X is not suitable for entry into the CODIS database for comparison purposes:*** The profile is too complex or not complete enough to be entered into the CODIS database.

***Reference samples will not be entered into CODIS:*** The Monroe County Crime Laboratory does not enter reference samples into CODIS unless submitted for the purpose of identifying a missing person.